



US006290956B1

(12) **United States Patent  
Bander**(10) **Patent No.: US 6,290,956 B1**  
(45) **Date of Patent: Sep. 18, 2001**(54) **TREATMENT AND DIAGNOSIS OF  
PROSTATE CANCER**(75) **Inventor:** Neil H. Bander, Chappaqua, NY (US)(73) **Assignee:** Cornell Research Foundation, Inc.,  
Ithaca, NY (US)(\*) **Notice:** Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.(21) **Appl. No.:** 09/357,710(22) **Filed:** Jul. 20, 1999**Related U.S. Application Data**(62) Division of application No. 08/838,682, filed on Apr. 9,  
1997, now Pat. No. 6,107,090.(60) Provisional application No. 60/016,976, filed on May 6,  
1996, and provisional application No. 60/022,125, filed on  
Jul. 18, 1996.(51) **Int. Cl.<sup>7</sup>** ..... A61K 39/395; A61K 39/40;  
A61K 39/42(52) **U.S. Cl.** ..... 424/130.1; 424/139.1;  
424/141.1; 424/138.1(58) **Field of Search** ..... 435/7.23, 7.1,  
435/7.2; 436/64; 530/387.1, 387.9, 588.8;  
424/130.1(56) **References Cited****U.S. PATENT DOCUMENTS**

4,814,275	3/1989	Durda et al.	435/240.27
4,863,851	9/1989	McEwan et al.	435/7
4,863,854	9/1989	Mattes et al.	435/7
5,013,645	5/1991	Kim	435/7.92
5,118,611	6/1992	Smith et al.	435/7.23
5,162,504	11/1992	Horoszewicz	530/388.2
5,227,471	7/1993	Wright, Jr.	530/388.8
5,229,289	7/1993	Kjeldsen et al.	435/240.27
5,314,996	5/1994	Wright, Jr.	530/387.3
5,342,924	8/1994	Chang	530/387.9
5,489,525	2/1996	Pastan	435/7.23
5,538,866	7/1996	Israeli et al.	435/69.3
5,599,677	2/1997	Dowell et al.	435/7.4
5,855,866	1/1999	Thorpe et al.	424/1.49
5,877,289	3/1999	Thorpe et al.	530/387.1

**OTHER PUBLICATIONS**Chang et al., "Five Different Anti-Prostate-specific Mem-  
brane Antigen (PSMA) Antibodies Confirm PSMA Expres-  
sion in Tumor-associated Neovasculture," *Cancer Res.*,  
59:3192-3198 (1999).Coleman et al., "Fundamental Immunology," Wm. C.  
Brown Publishers, p. 76 (1989).Harlow et al., "Antibodies: A laboratory manual", Cold  
Spring Harbor Laboratory, pp. 139-243 (1988).Heston, "Characterization and Glutamyl Preferring Carbox-  
ypeptidase Function of Prostate Specific Membrane Anti-  
gen: A Novel Folate Hydrolase" (*Urology* 49 (Supp. 3A):  
104-112; 106 (1997).Horoszewicz et al. "Monoclonal Antibodies to a New Anti-  
genic Marker in Epithelial Prostatic Cells and Serum of  
Prostatic Cancer Patients," *Anticancer Research*, 7:927-936  
(1987).Israeli et al., "Expression of the Prostate-Specific Mem-  
brane Antigen", *Cancer Research*, 54:1807-1811 (1994).Israeli et al., "Molecular Cloning of a Complementary DNA  
Encoding a Prostate-Specific Membrane Antigen", *Cancer  
Research*, 53:227-230 (1993).Jain, R.K. "Vascular and interstitial barriers to delivery of  
therapeutic agents in tumors" *Cancer and Metastasis  
Reviews*, vol. 9, pp. 253-266 (1990).Liu et al., "Constitutive and Antibody-Induced Internaliza-  
tion of Prostate-Specific Membrane Antigen," *Cancer Res.*  
58:4055-60 (1998).Liu, et. al., "Monoclonal Antibodies to the Extracellular  
Domain of Prostate-specific Membrane Antigen also React  
with Tumor Vascular Endothelium," *Cancer Res.*  
57:3629-34 (1997).Murphy et al., "Measurement of Prostate-Specific Mem-  
brane Antigen in the Serum With a New Antibody," *The  
Prostate*, 28:266-271 (1996).Rochon et al., "Western Blot Assay for Prostate-Specific  
Membrane Antigen in Serum of Prostate Cancer Patients,"  
*The Prostate*, 25:219-223 (1994).Schlom, "Monoclonal antibodies: They're more and less  
than you think", in *Molecular Foundations of Oncology*, ed.  
by Broder, William & Wilkins (1991).Silver et al., "Prostate-specific Membrane Antigen Expres-  
sion in Normal and Malignant Human Tissues," *Clin. Can-  
cer Res.* 3:81-85 (1997).Troyer et al., "Biochemical Characterization and Mapping  
of the 7E11-C5.3 Epitope of the Prostate-Specific Mem-  
brane Antigen," *Urol. Oncol.*, 1:29-37 (1995).Troyer et al., "Detection and Characterization of the Pros-  
tate-Specific Membrane Antigen (PSMA) in Tissue Extracts  
and Body Fluids," *Int. J. Chancer*, 62:552-558 (1995).Troyer et al., "Location of prostate-specific membrane anti-  
gen in the LNCaP prostate carcinoma cell line," *Prostate*,  
30:232-242. (1997).

(List continued on next page.)

**Primary Examiner**—Susan Ungar**Assistant Examiner**—Gary B. Nickol(74) **Attorney, Agent, or Firm**—Lyon & Lyon LLP

(57)

**ABSTRACT**

The present invention is directed to the use of antibodies or binding portions thereof, probes, ligands, or other biological agents which either recognize an extracellular domain of prostate specific membrane antigen or bind to and are internalized with prostate specific membrane antigen. These biological agents can be labeled and used for detection of normal, benign hyperplastic, and cancerous prostate epithelial cells or portions thereof. They also can be used alone or bound to a substance effective to ablate or kill such cells as a therapy for prostate cancer. Also disclosed are four hybridoma cell lines, each of which produces a monoclonal antibody recognizing extracellular domains of prostate specific membrane antigens of normal, benign hyperplastic, and cancerous prostate epithelial cells or portions thereof.

**3 Claims, 11 Drawing Sheets**

Applicants: Ron S. Israeli, et al.

Serial No. : 08/470,735

Filed: June 6, 1995


Exhibit I

OTHER PUBLICATIONS


Wright et al., "Expression of Prostate-Specific Membrane Antigen in Normal, Benign, and Malignant Prostate Tissues," *Urol Oncol.*, 1:18-28 (1995).

Yang et al., AACR Abstract #2996 (1998); "Alpha particle emitter therapy of micrometastases:  $^{213}\text{Bi}$ -J5 . . . (anti-P-SMA) treatment of LnCaP spheroids," Proceedings of the American Association for Cancer Research, 39:440 (1998).


---



**FIG. 1**




**FIG. 2**




**FIG. 3**



**FIG. 4**



**FIG. 5**

**FIG. 6**



ENZYMES: ALL 74 ENZYMES (NO FILTER)

SETTINGS: LINEAR, CERTAIN SITES ONLY, STANDARD GENETIC CODE




FIG. 7

**FIG. 8**




FIG. 9

ENZYMES: ALL 74 ENZYMES (NO FILTER):  
 SETTINGS: LINEAR, CERTAIN SITES ONLY, STANDARD GENETIC CODE




FIG. 10

## LIPMAN-PEARSON PROTEIN ALIGNMENT

KTUPLE: 2; GAP PENALTY: 4; GAP LENGTH PENALTY: 12

SEQ1(1>107)	SEQ2(1 > 111)	SIMILARITY	GAP	GAP	CONSENSUS
J591VK.PRO	MUVKV.PRO	INDEX	NUMBER	LENGTH	LENGTH
(1>107)	(1>109)	60.4	2	2	109

↓10      ↓20      ↓30      ↓40      ↓50  
NIVMTQSPKSMMSVGERVILTCKAS-ENVVTYVSWYQOKPEQSPKLLITYGASNRYTGVP  
:I MTQSP.S:S S:G:RVT:TC:AS ::: .Y::WYQOKP. SPKLLIY AS. .:GVP  
DIQMTQSPSSLSASLGDRVTTTCRASQDDISNYLAWYQOKPGGSPKLLIYYASRLHSGVP  
↑10      ↑20      ↑30      ↑40      ↑50      ↑60  
↓60      ↓70      ↓80      ↓90      ↓100  
DRFTGSGSATDFTLTISVQAEADLADYHCGQGY-SY-PYTFGGGKLEIK  
.RF:GSGS:TD::LTIS:::ED:A.Y C QG : P TFGGGKLEIK  
SRFSGSGSGIDYSLTISNLEQEDIATYFCQCGNLPPTFTFGGGKLEIK  
↑70      ↑80      ↑90      ↑100

FIG. 11

## TREATMENT AND DIAGNOSIS OF PROSTATE CANCER

This application is a divisional of application Ser. No. 08/838,682, filed on Apr. 9, 1997 now U.S. Pat. No. 6,107,090; which claims the benefit of U.S. Provisional Patent Application Ser. No. 60/016,976, filed May 6, 1996, now abandoned and U.S. Provisional Patent Application Ser. No. 60/022,125, filed Jul. 18, 1996 now abandoned.

### FIELD OF THE INVENTION

The present invention relates to the treatment and diagnosis of prostate cancer with biological agents.

### BACKGROUND OF THE INVENTION

Prostate cancer is the most common cancer in men with an estimated 317,000 cases in 1996 in the United States. It is the second leading cause of death among men who die from neoplasia with an estimated 40,000 deaths per year. Prompt detection and treatment is needed to limit mortality caused by prostate cancer.

#### Detection of Prostate Cancer

When it metastasizes, prostatic cancer has a distinct predilection for bone and lymph nodes. Saitoh et al., "Metastatic Patterns of Prostatic Cancer. Correlation Between Sites And Number Of Organs Involved," *Cancer*, 54:3078-3084 (1984). At the time of clinical diagnosis, as many as 25% of patients have bone metastasis demonstrable by radionuclide scans. Murphy, G. P., et al., "The National Survey Of Prostate Cancer In The United States By The American College Of Surgeons," *J. Urol.*, 127:928-939 (1982). Accurate clinical evaluation of nodal involvement has proven to be difficult. Imaging techniques such as computed tomography ("CT") or magnetic resonance ("MR") imaging are unable to distinguish metastatic prostate cancer involvement of lymph nodes by criterion other than size (i.e., >1 cm). Therefore, by definition, these imaging modalities are inherently insensitive in the detection of small volume (<1 cm) disease as well as non-specific in the detection of larger volume adenopathy. A recent study assessed the accuracy of MR in patients with clinically localized prostate cancer. Rifkin et al., "Comparison Of Magnetic Resonance Imaging And Ultrasonography In Staging Early Prostate Cancer," *N. Engl. J. Med.*, 323:621-626 (1990). In this study, 194 patients underwent an MR and 185 of these patients had a lymph node dissection. 23 (13%) patients had pathologically involved lymph nodes. MR was suspicious in only 1 of these 23 cases resulting in a sensitivity of 4%. Similar results have also been noted with CT scans. Gasser et al., "MRI And Ultrasonography In Staging Prostate Cancer," *N. Engl. J. Med. (Correspondence)*, 324 (7):49-495 (1991).

The elevation of serum acid phosphatase activity in patients having metastasized prostate carcinoma was first reported by Gutman et al., *J. Clin. Invest.* 17:473 (1938). In cancer of the prostate, prostatic acid phosphatase is released from the cancer tissue into the blood stream with the result that the total serum acid phosphatase level can be greatly increased above normal values. Numerous studies of this enzyme and its relation to prostatic cancer have been made since that time, e.g. Yam, *Amer. J. Med.* 56:604 (1974). However, the measurement of serum acid phosphatase is elevated in about 65-90 percent of patients having carcinoma of the prostate with bone metastasis; in about 30 percent of patients without roentgenological evidence of bone metastasis; and in about only 5-10 percent of patients lacking clinically demonstrable metastasis.

Prior art attempts to develop a specific test for prostatic acid phosphatase have met with only limited success, because techniques which rely on enzyme activity on a so-called "specific" substrate cannot take into account other biochemical and immunochemical differences among the many acid phosphatases which are unrelated to enzyme activity of prostate origin. In the case of isoenzymes, i.e. genetically defined enzymes having the same characteristic enzyme activity and a similar molecular structure but differing in amino acid sequences and/or content and, therefore, immunochemically distinguishable, it would appear inherently impossible to distinguish different isoenzyme forms merely by the choice of a particular substrate. It is, therefore, not surprising that none of these prior art methods is highly specific for the direct determination of prostatic acid phosphatase activity; e.g. see *Cancer* 5:236 (1952); *J. Lab. Clin. Med.* 82:486 (1973); *Clin. Chem. Acta.* 44:21 (1973); and *J. Physiol. Chem.* 356:1775 (1975).

In addition to the aforementioned problems of non-specificity which appear to be inherent in many of the prior art reagents employed for the detection of prostate acid phosphatase, there have been reports of elevated serum acid phosphatase associated with other diseases, which further complicates the problem of obtaining an accurate clinical diagnosis of prostatic cancer. For example, Tuchman et al., *Am. J. Med.* 27:959 (1959) noted that serum acid phosphatase levels appear to be elevated in patients with Gaucher's disease.

Due to the inherent difficulties in developing a "specific" substrate for prostate acid phosphatase, several researchers have developed immunochemical methods for the detection of prostate acid phosphatase. However, the previously reported immunochemical methods have drawbacks of their own which have precluded their widespread acceptance. For example, Shulman et al., *Immunology* 93:474 (1964) described an immuno-diffusion test for the detection of human prostate acid phosphatase. Using antisera prepared from a prostatic fluid antigen obtained by rectal massage from patients with prostatic disease, no cross-reactivity precipitin line was observed in the double diffusion technique against extracts of normal kidney, testicle, liver, and lung. However, this method has the disadvantages of limited sensitivity, even with the large amounts of antigen employed, and of employing antisera which may cross-react with other, antigenically unrelated serum protein components present in prostatic fluid.

WO 79/00475 to Chu et. al. describes a method for the detection of prostatic acid phosphatase isoenzyme patterns associated with prostatic cancer which obviates many of the above drawbacks. However, practical problems are posed by the need for a source of cancerous prostate tissue from which the diagnostically relevant prostatic acid phosphatase isoenzyme patterns associated with prostatic cancer are extracted for the preparation of antibodies thereto.

In recent years, considerable effort has been spent to identify enzyme or antigen markers for various types of malignancies with the view towards developing specific diagnostic reagents. The ideal tumor marker would exhibit, among other characteristics, tissue or cell-type specificity. Previous investigators have demonstrated the occurrence of human prostate tissue-specific antigens.

#### Treatment of Prostate Cancer

As described in W. J. Catalona, "Management of Cancer of the Prostate," *New Engl. J. Med.*, 331(15):996-1004 (1994), the management of prostate cancer can be achieved by watchful waiting, curative treatment, and palliation.

For men with a life expectancy of less than 10 years, watchful waiting is appropriate where low-grade, low-stage

prostate cancer is discovered at the time of a partial prostatectomy for benign hyperplasia. Such cancers rarely progress during the first five years after detection. On the other hand, for younger men, curative treatment is often more appropriate.

Where prostate cancer is localized and the patient's life expectancy is 10 years or more, radical prostatectomy offers the best chance for eradication of the disease. Historically, the drawback of this procedure is that most cancers had spread beyond the bounds of the operation by the time they were detected. However, the use of prostate-specific antigen testing has permitted early detection of prostate cancer. As a result, surgery is less extensive with fewer complications. Patients with bulky, high-grade tumors are less likely to be successfully treated by radical prostatectomy.

After surgery, if there are detectable serum prostate-specific antigen concentrations, persistent cancer is indicated. In many cases, prostate-specific antigen concentrations can be reduced by radiation treatment. However, this concentration often increases again within two years.

Radiation therapy has also been widely used as an alternative to radical prostatectomy. Patients generally treated by radiation therapy are those who are older and less healthy and those with higher-grade, more clinically advanced tumors. Particularly preferred procedures are external-beam therapy which involves three dimensional, conformal radiation therapy where the field of radiation is designed to conform to the volume of tissue treated; interstitial-radiation therapy where seeds of radioactive compounds are implanted using ultrasound guidance; and a combination of external-beam therapy and interstitial-radiation therapy.

For treatment of patients with locally advanced disease, hormonal therapy before or following radical prostatectomy or radiation therapy has been utilized. Hormonal therapy is the main form of treating men with disseminated prostate cancer. Orchiectomy reduces serum testosterone concentrations, while estrogen treatment is similarly beneficial. Diethylstilbestrol from estrogen is another useful hormonal therapy which has a disadvantage of causing cardiovascular toxicity. When gonadotropin-releasing hormone agonists are administered testosterone concentrations are ultimately reduced. Flutamide and other nonsteroidal, anti-androgen agents block binding of testosterone to its intracellular receptors. As a result, it blocks the effect of testosterone, increasing serum testosterone concentrations and allows patients to remain potent—a significant problem after radical prostatectomy and radiation treatments.

Cytotoxic chemotherapy is largely ineffective in treating prostate cancer. Its toxicity makes such therapy unsuitable for elderly patients. In addition, prostate cancer is relatively resistant to cytotoxic agents.

#### Use of Monoclonal Antibodies in Prostate Cancer Detection and Treatment

Theoretically, radiolabeled monoclonal antibodies ("mAbs") offer the potential to enhance both the sensitivity and specificity of detecting prostatic cancer within lymph nodes and elsewhere. While many mAbs have previously been prepared against prostate related antigens, none of these mAbs were specifically generated with an imaging objective in mind. Nevertheless, the clinical need has led to evaluation of some of these mAbs as possible imaging agents. Vihko et al., "Radioimaging of Prostatic Carcinoma With Prostatic Acid Phosphatase—Specific Antibodies," *Biotechnology in Diagnostics*, 131–134 (1985); Babaian et al., "Radioimmunological Imaging of Metastatic Prostatic Cancer With 111-Indium-Labeled Monoclonal Antibody PAY 276," *J. Urol.*, 137:439–443 (1987); Leroy et al.,

"Radioimmunodetection Of Lymph Node Invasion In Prostatic Cancer. The Use Of Iodine 123 (123-I)-Labeled Monoclonal Anti-Prostatic Acid Phosphatase (PAP) 227 A F (ab')<sub>2</sub> Antibody Fragments In Vivo," *Cancer*, 64:1–5 (1989); Meyers et al., "Development Of Monoclonal Antibody Imaging Of Metastatic Prostatic Carcinoma," *The Prostate*, 14:209–220 (1989).

In some cases, the monoclonal antibodies developed for detection and/or treatment of prostate cancer recognize antigens specific to malignant prostatic tissues. Such antibodies are thus used to distinguish malignant prostatic tissue (for treatment or detection) from benign prostatic tissue. See U.S. Pat. No. 4,970,299 to Bazinet et al. and U.S. Pat. No. 4,902,615 to Freeman et al.

Other monoclonal antibodies react with surface antigens on all prostate epithelial cells whether cancerous or benign. See U.S. Pat. Nos. 4,446,122 and Re 33,405 to Chu et al., U.S. Pat. No. 4,863,851 to McEwan et al., and U.S. Pat. No. 5,055,404 to Ueda et al. However, the antigens detected by these monoclonal antibodies are present in the blood and, therefore, compete with antigens at tumor sites for the monoclonal antibodies. This causes background noise which makes the use of such antibodies inadequate for in vivo imaging. In therapy, such antibodies, if bound to a cytotoxic agent, could be harmful to other organs.

Horoszewicz et al., "Monoclonal Antibodies to a New Antigenic Marker in Epithelial Prostatic Cells and Serum of Prostatic Cancer Patients," *Anticancer Research*, 7:927–936 (1987) ("Horoszewicz") and U.S. Pat. No. 5,162,504 to Horoszewicz describe an antibody, designated 7E11, which recognizes prostate specific membrane antigen ("PSMA"). Israeli et al., "Molecular Cloning of a Complementary DNA Encoding a Prostate-specific Membrane Antigen," *Cancer Research*, 53:227–230 (1993) ("Israeli") describes the cloning and sequencing of PSMA and reports that PSMA is prostate-specific and shows increased expression levels in metastatic sites and in hormone-refractory states. Other studies have indicated that PSMA is more strongly expressed in prostate cancer cells relative to cells from the normal prostate or from a prostate with benign hyperplasia. Furthermore, PSMA is not found in serum (Troyer et al., "Detection and Characterization of the Prostate-Specific Membrane Antigen (PSMA) in Tissue Extracts and Body Fluids," *Int. J. Cancer*, 62:552–558 (1995)).

These characteristics make PSMA an attractive target for antibody mediated targeting for imaging and therapy of prostate cancer. Imaging studies using indium-labeled 7E11 have indicated that the antibody localizes quite well to both the prostate and to sites of metastasis. In addition, 7E11 appears to have clearly improved sensitivity for detecting lesions compared to other currently available imaging techniques, such as CT and MR imaging or bone scan. Bander, "Current Status of Monoclonal Antibodies for Imaging and Therapy of Prostate Cancer," *Sem. In Oncology*, 21:607–612 (1994).

However, the use of 7E11 and other known antibodies to PSMA to mediate imaging and therapy has several disadvantages. First, PSMA is an integral membrane protein known to have a short intracellular tail and a long extracellular domain. Biochemical characterization and mapping (Troyer et al., "Biochemical Characterization and Mapping of the 7E11-C5.3 Epitope of the Prostate-specific Membrane Antigen," *Urol. Oncol.*, 1:29–37 (1995)) have shown that the epitope or antigenic site to which the 7E11 antibody binds is present on the intracellular portion of the molecule. Because antibody molecules do not, under normal circumstances, cross the cell membrane unless they bind to

the extracellular portion of a molecule and become translocated intracellularly, the 7E11 antibody does not have access to its antigenic target site in an otherwise healthy, viable cell.

Consequently, imaging using 7E11 is limited to the detection of dead cells within tumor deposits. Additionally, the therapeutic use of the 7E11 antibody is limited, because only cells that are already dead or tissue containing a large proportion of dead cells can be effectively targeted.

The present invention is directed to overcoming the deficiencies of prior art antibodies in diagnosing and treating prostate cancer.

#### SUMMARY OF THE INVENTION

One aspect of the present invention relates to a method of ablating or killing normal, benign hyperplastic, and cancerous prostate epithelial cells. The process involves providing a biological agent which recognizes an extracellular domain of prostate specific membrane antigen. The biological agent can be used alone or can be bound to a substance effective to kill the cells upon binding of the biological agent to the cells. These biological agents are then contacted with the cells under conditions effective to permit both binding of the biological agent to the extracellular domain of the prostate specific membrane antigen and killing or ablating of the cells.

In another particularly preferred embodiment of the method of ablating or killing normal, benign hyperplastic, and cancerous prostate epithelial cells in accordance with the present invention, the biological agent binds to and is internalized with the prostate specific membrane antigen of such cells. Preferred biological agents for use in the method of ablating or killing normal, benign hyperplastic, and cancerous prostate epithelial cells in accordance with the present invention are antibodies or binding portions thereof, probes, or ligands.

Another aspect of the present invention relates to a method of detecting normal, benign hyperplastic, and cancerous prostate epithelial cells or portions thereof in a biological sample. This method involves providing a biological agent which binds to an extracellular domain of prostate specific membrane antigen. The biological agent is bound to a label effective to permit detection of the cells or portions thereof upon binding of the biological agent to the cells or portions thereof. The biological sample is contacted with the biological agent having a label under conditions effective to permit binding of the biological agent to the extracellular domain of the prostate specific membrane antigen of any of the cells or portions thereof in the biological sample. The presence of any cells or portions thereof in the biological sample is detected by detection of the label.

In a particularly preferred embodiment of the method of detecting normal, benign hyperplastic, and cancerous prostate epithelial cells in accordance with the present invention, the biological agent binds to and is internalized with the prostate specific membrane antigen of such cells. Preferred biological agents for use in the method of detecting normal, benign hyperplastic, and cancerous prostate epithelial cells in accordance with the present invention are antibodies or binding portions thereof, probes, or ligands.

Another aspect of the present invention pertains to a biological agent that recognizes an extracellular domain of prostate specific membrane antigen. In a preferred embodiment, the isolated biological agent binds to and is internalized with the prostate specific membrane antigen. Preferred isolated biological agents which recognize an

extracellular domain of prostate specific membrane antigen in accordance with the present invention are isolated antibodies or binding portions thereof, probes, or ligands. Hybridoma cell lines that produce monoclonal antibodies of these types are also disclosed.

The biological agents of the present invention recognize the extracellular domain of antigens of normal, benign hyperplastic, and cancerous prostate epithelial cells. Unlike the 7E11 antibody, which recognizes an epitope of prostate-associated antigens which are exposed extracellularly only after cell lysis, the biological agents of the present invention bind to antigenic epitopes which are extracellularly exposed in living prostate cells. Using the biological agents of the present invention, living, unfixed normal, benign hyperplastic, and cancerous prostate epithelial cells can be targeted, which makes treatment and diagnosis more effective. In a preferred embodiment, the biological agents of the present invention also bind to and are internalized with the prostate specific membrane antigen, which permits the therapeutic use of intracellularly acting cytotoxic agents.

#### BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is an immuno-electron micrograph of gold-labeled monoclonal antibody J591 on the surface of LNCaP cells prior to incubation.

FIG. 2 is an immuno-electron micrograph of gold-labeled monoclonal antibody J591 after 5 minutes incubation at 37° C. LNCaP cells.

FIG. 3 is an immuno-electron micrograph of gold-labeled monoclonal antibody J591 after 10 minutes incubation at 37° C. LNCaP cells.

FIG. 4 is an immuno-electron micrograph of gold-labeled monoclonal antibody J591 after 15 minutes incubation at 37° C. LNCaP cells.

FIG. 5 is an immuno-electron micrograph of gold-labeled monoclonal antibody J591 after 15 minutes at 37° C. showing J591 within endosomes.

FIG. 6 summarizes the sequencing strategy of the heavy chain of monoclonal antibody J591.

FIG. 7 shows the nucleotide sequence of the heavy chain of monoclonal antibody J591 (designated SEQ. ID. No. 1), the nucleotide sequence of the corresponding reverse, non-coding strand (designated SEQ. ID. No. 2), and the corresponding deduced amino acid sequences (designated SEQ. ID. Nos. 3, 4, and 5).

FIG. 8 is a comparison of the heavy chain of monoclonal antibody J591 with the consensus sequence for Mouse Heavy Chains Subgroup IIA (designated SEQ ID No. 20). As conventionally recognized, the identity of consensus between the two sequences is displayed in the line between the top J591 sequence (designated SEQ ID No. 8) and the bottom sequence of mouse antibody heavy chain subgroup IIA (designated SEQ ID No. 20).

FIG. 9 summarizes the sequencing strategy of the kappa light chain of monoclonal antibody J591.

FIG. 10 shows the nucleotide sequences of the kappa light chain of monoclonal antibody J591 (designated SEQ. ID. No. 9), the nucleotide sequence of the corresponding reverse, non-coding strand (designated SEQ. ID. No. 10), and the corresponding deduced amino acid sequence (designated SEQ. ID. Nos. 11, 12, and 13).

FIG. 11 is a comparison of the kappa light chain of monoclonal antibody J591 with the consensus sequence for Mouse Kappa Chains Subgroup V (designated SEQ ID No. 21). As conventionally recognized, the identity of consensus



between the two sequences is displayed in the line between the top J591 sequence (designated SEQ ID No. 19) and the bottom sequence of mouse antibody kappa light chain subgroup V (designated SEQ ID No. 21).

#### DETAILED DESCRIPTION OF THE INVENTION

One aspect of the present invention relates to a method of ablating or killing normal, benign hyperplastic, and cancerous prostate epithelial cells. The process involves providing a biological agent, such as an antibody or binding portion thereof, probe, or ligand, which binds to an extracellular domain of prostate specific membrane antigen of (i.e., a portion of prostate specific membrane antigen which is external to) such cells. The biological agent can be used alone or can be bound to a substance effective to kill the cells upon binding of the biological agent to the cells. These biological agents are then contacted with the cells under conditions effective to permit both binding of the biological agent to the extracellular domain of the prostate specific membrane antigen and killing or ablating of the cells. In its preferred form, such contacting is carried out in a living mammal by administering the biological agent to the mammal under conditions effective to permit both binding of the biological agent to the extracellular domain of the prostate specific membrane antigen and killing or ablating of the cells. Such administration can be carried out orally or parenterally.

In a particularly preferred embodiment of the method of ablating or killing normal, benign hyperplastic, and cancerous prostate epithelial cells in accordance with the present invention, the biological agent binds to and is internalized with the prostate specific membrane antigen of such cells. Again, the biological agent can be used alone. Alternatively, the biological agent can be bound to a substance effective to kill the cells upon binding of the biological agent to prostate specific membrane antigen and upon internalization of the biological agent with the prostate specific membrane antigen.

The mechanism by which the biological agent is internalized with the prostate specific membrane antigen is not critical to the practice of the present invention. For example, the biological agent can induce internalization of the prostate specific membrane antigen. Alternatively, internalization of the biological agent can be the result of routine internalization of prostate specific membrane antigen.

Another aspect of the present invention relates to a method of detecting normal, benign hyperplastic, and cancerous epithelial cells or portions thereof in a biological sample. This method involves providing a biological agent, such as an antibody or binding portion thereof, probe, or ligand, which binds to an extracellular domain of prostate specific membrane antigen of such cells. The biological agent is bound to a label effective to permit detection of the cells or portions (e.g., prostate specific membrane antigen or fragments thereof liberated from such normal, benign hyperplastic, and cancerous cells) thereof upon binding of the biological agent to the cells or portions thereof. The biological sample is contacted with the biological agent having a label under conditions effective to permit binding of the biological agent to the extracellular domain of the prostate specific membrane antigen of any of the cells or portions thereof in the biological sample. The presence of any cells or portions thereof in the biological sample is detected by detection of the label. In its preferred form, such contacting is carried out in a living mammal and involves

administering the biological agent to the mammal under conditions effective to permit binding of the biological agent to the prostate specific membrane antigen of any of the cells or portions thereof in the biological sample. Again, such administration can be carried out orally or parenterally.

The method of the present invention can be used to screen patients for diseases associated with the presence of normal, benign hyperplastic, and cancerous epithelial cells or portions thereof. Alternatively, it can be used to identify the recurrence of such diseases, particularly when the disease is localized in a particular biological material of the patient. For example, recurrence of prostatic disease in the prostatic fossa may be encountered following radical prostatectomy. Using the method of the present invention, this recurrence can be detected by administering a short range radiolabeled antibody to the mammal and then detecting the label rectally, such as with a transrectal detector probe.

Alternatively, the contacting step can be carried out in a sample of serum or urine or other body fluids, such as to detect the presence of PSMA in the body fluid. When the contacting is carried out in a serum or urine sample, it is preferred that the biological agent recognize substantially no antigens circulating in the blood other than PSMA. Since intact prostate cells do not excrete or secrete PSMA into the extracellular environment, detecting PSMA in serum, urine, or other body fluids generally indicates that prostate cells are being lysed. Thus, the biological agents and methods of the present invention can be used to determine the effectiveness of a prostate cancer treatment protocol by monitoring the level of PSMA in serum, urine or other body fluids.

In a particularly preferred embodiment of the method of detecting normal, benign hyperplastic, and cancerous prostate epithelial cells in accordance with the present invention, the biological agent, such as the antibody or binding portion thereof, probe, or ligand, binds to and is internalized with the prostate specific membrane antigen of such cells. Again, the biological agent is bound to a label effective to permit detection of the cells or portions thereof upon binding of the biological agent to and internalization of the biological agent with the prostate specific membrane antigen.

As indicated above, biological agents suitable for either killing, ablating, or detecting normal, benign hyperplastic, and cancerous prostate epithelial cells include antibodies, such as monoclonal or polyclonal antibodies. In addition, antibody fragments, half-antibodies, hybrid derivatives, probes, and other molecular constructs may be utilized. These biological agents, such as antibodies, binding portions thereof, probes, or ligands, bind to extracellular domains of prostate specific membrane antigens or portions thereof in normal, benign hyperplastic, and cancerous prostate epithelial cells. As a result, the biological agents bind to all such cells, not only to cells which are fixed or cells whose intracellular antigenic domains are otherwise exposed to the extracellular environment. Consequently, binding of the biological agents is concentrated in areas where there are prostate epithelial cells, irrespective of whether these cells are fixed or unfixed, viable or necrotic. Additionally or alternatively, these biological agents, such as antibodies, binding portions thereof, probes, or ligands, bind to and are internalized with prostate specific membrane antigens or portions thereof in normal, benign hyperplastic, and cancerous prostate epithelial cells.

Monoclonal antibody production may be effected by techniques which are well-known in the art. Basically, the process involves first obtaining immune cells (lymphocytes) from the spleen of a mammal (e.g., mouse) which has been

previously immunized with the antigen of interest either in vivo or in vitro. The antibody-secreting lymphocytes are then fused with (mouse) myeloma cells or transformed cells, which are capable of replicating indefinitely in cell culture, thereby producing an immortal, immunoglobulin-secreting cell line. The resulting fused cells, or hybridomas, are cultured, and the resulting colonies screened for the production of the desired monoclonal antibodies. Colonies producing such antibodies are cloned, and grown either in vivo or in vitro to produce large quantities of antibody. A description of the theoretical basis and practical methodology of fusing such cells is set forth in Kohler and Milstein, *Nature* 256:495 (1975), which is hereby incorporated by reference.

Mammalian lymphocytes are immunized by in vivo immunization of the animal (e.g., a mouse) with the protein or polypeptide of the present invention. Such immunizations are repeated as necessary at intervals of up to several weeks to obtain a sufficient titer of antibodies. Following the last antigen boost, the animals are sacrificed and spleen cells removed.

Fusion with mammalian myeloma cells or other fusion partners capable of replicating indefinitely in cell culture is effected by standard and well-known techniques, for example, by using polyethylene glycol ("PEG") or other fusing agents (See Milstein and Kohler, *Eur. J. Immunol.* 6:511 (1976), which is hereby incorporated by reference). This immortal cell line, which is preferably murine, but may also be derived from cells of other mammalian species, including but not limited to rats and humans, is selected to be deficient in enzymes necessary for the utilization of certain nutrients, to be capable of rapid growth, and to have good fusion capability. Many such cell lines are known to those skilled in the art, and others are regularly described.

Procedures for raising polyclonal antibodies are also well known. Typically, such antibodies can be raised by administering the protein or polypeptide of the present invention subcutaneously to New Zealand white rabbits which have first been bled to obtain pre-immune serum. The antigens can be injected at a total volume of 100  $\mu$ l per site at six different sites. Each injected material will contain synthetic surfactant adjuvant pluronic polyols, or pulverized acrylamide gel containing the protein or polypeptide after SDS-polyacrylamide gel electrophoresis. The rabbits are then bled two weeks after the first injection and periodically boosted with the same antigen three times every six weeks. A sample of serum is then collected 10 days after each boost. Polyclonal antibodies are then recovered from the serum by affinity chromatography using the corresponding antigen to capture the antibody. Ultimately, the rabbits are euthenized with pentobarbital 150 mg/Kg IV. This and other procedures for raising polyclonal antibodies are disclosed in E. Harlow, et. al., editors, *Antibodies: A Laboratory Manual* (1988), which is hereby incorporated by reference.

In addition to utilizing whole antibodies, the processes of the present invention encompass use of binding portions of such antibodies. Such binding portions include Fab fragments, F(ab')<sub>2</sub> fragments, and Fv fragments. These antibody fragments can be made by conventional procedures, such as proteolytic fragmentation procedures, as described in J. Goding, *Monoclonal Antibodies: Principles and Practice*, pp. 98-118 (N.Y. Academic Press 1983), which is hereby incorporated by reference.

Alternatively, the processes of the present invention can utilize probes or ligands found either in nature or prepared synthetically by recombinant DNA procedures or other biological or molecular procedures. Suitable probes or

ligands are molecules which bind to the extracellular domains of prostate specific membrane antigens identified by the monoclonal antibodies of the present invention. Other suitable probes or ligands are molecules which bind to and are internalized with prostate specific membrane antigens. Such probes or ligands can be, for example, proteins, peptides, lectins, or nucleic acid probes.

It is particularly preferred to use the monoclonal antibodies identified below in Table 1.

TABLE 1

Monoclonal Antibody Name	ATCC Designation for Hybridoma Cell Line
E99	HB-12101
J415	HB-12109
J533	HB-12127
J591	HB-12126

These antibodies can be used alone or as a component in a mixture with other antibodies or other biological agents to treat or image prostate epithelial cells with varying surface antigen characteristics.

Regardless of whether the biological agents are used for treatment or therapy, they can be administered orally, parenterally, subcutaneously, intravenously, intramuscularly, intraperitoneally, by intranasal instillation, by intracavitary or intravesical instillation, intraocularly, intraarterially, intralesionally, or by application to mucous membranes, such as, that of the nose, throat, and bronchial tubes. They may be administered alone or with pharmaceutically or physiologically acceptable carriers, excipients, or stabilizers, and can be in solid or liquid form such as, tablets, capsules, powders, solutions, suspensions, or emulsions.

The solid unit dosage forms can be of the conventional type. The solid form can be a capsule, such as an ordinary gelatin type containing the biological agent, such as an antibody or binding portion thereof, of the present invention and a carrier, for example, lubricants and inert fillers such as, lactose, sucrose, or cornstarch. In another embodiment, these compounds are tableted with conventional tablet bases such as lactose, sucrose, or cornstarch in combination with binders like acacia, cornstarch, or gelatin, disintegrating agents such as, cornstarch, potato starch, or alginic acid, and a lubricant like stearic acid or magnesium stearate.

The biological agent of the present invention may also be administered in injectable dosages by solution or suspension of these materials in a physiologically acceptable diluent with a pharmaceutical carrier. Such carriers include sterile liquids such as water and oils, with or without the addition of a surfactant and other pharmaceutically and physiologically acceptable carrier, including adjuvants, excipients or stabilizers. Illustrative oils are those of petroleum, animal, vegetable, or synthetic origin, for example, peanut oil, soybean oil, or mineral oil. In general, water, saline, aqueous dextrose and related sugar solution, and glycols such as, propylene glycol or polyethylene glycol, are preferred liquid carriers, particularly for injectable solutions.

For use as aerosols, the biological agent of the present invention in solution or suspension may be packaged in a pressurized aerosol container together with suitable propellants, for example, hydrocarbon propellants like propane, butane, or isobutane with conventional adjuvants. The materials of the present invention also may be administered in a non-pressurized form such as in a nebulizer or atomizer.

The biological agents may be utilized to detect normal, benign hyperplastic, and cancerous prostate epithelial cells in vivo. This is achieved by labeling the biological agent, administering the labeled biological agent to a mammal, and then imaging the mammal.

Examples of labels useful for diagnostic imaging in accordance with the present invention are radiolabels such as  $^{131}\text{I}$ ,  $^{111}\text{In}$ ,  $^{123}\text{I}$ ,  $^{99\text{m}}\text{Tc}$ ,  $^{32}\text{P}$ ,  $^{125}\text{I}$ ,  $^3\text{H}$ ,  $^{14}\text{C}$ , and  $^{188}\text{Re}$ , fluorescent labels such as fluorescein and rhodamine, nuclear magnetic resonance active labels, positron emitting isotopes detectable by a positron emission tomography ("PET") scanner, chemiluminescers such as luciferin, and enzymatic markers such as peroxidase or phosphatase. Short-range radiation emitters, such as isotopes detectable by short-range detector probes, such as a transrectal probe, can also be employed. These isotopes and transrectal detector probes, when used in combination, are especially useful in detecting prostatic fossa recurrences and pelvic nodal disease. The biological agent can be labeled with such reagents using techniques known in the art. For example, see Wensel and Meares, *Radioimmunoimaging and Radioimmunotherapy*, Elsevier, New York (1983), which is hereby incorporated by reference, for techniques relating to the radiolabeling of antibodies. See also, D. Colcher et al., "Use of Monoclonal Antibodies as Radiopharmaceuticals for the Localization of Human Carcinoma Xenografts in Athymic Mice", *Meth. Enzymol.* 121: 802-816 (1986), which is hereby incorporated by reference.

A radiolabeled biological agent of this invention can be used for in vitro diagnostic tests. The specific activity of a tagged biological agent, such as a tagged antibody, binding portion thereof, probe, or ligand, depends upon the half-life, the isotopic purity of the radioactive label, and how the label is incorporated into the biological agent. Table 2 lists several commonly-used isotopes, their specific activities and half-lives. In immunoassay tests, the higher the specific activity, in general, the better the sensitivity.

TABLE 2

Isotope	Specific Activity of Pure Isotope (Curies/mole)	Half-Life
$^{14}\text{C}$	$6.25 \times 10^1$	5720 years
$^3\text{H}$	$2.01 \times 10^4$	12.5 years
$^{35}\text{S}$	$1.50 \times 10^6$	87 days
$^{125}\text{I}$	$2.18 \times 10^6$	60 days
$^{32}\text{P}$	$3.16 \times 10^6$	14.3 days
$^{131}\text{I}$	$1.62 \times 10^7$	8.1 days

Procedures for labeling biological agents with the radioactive isotopes listed in Table 2 are generally known in the art. Tritium labeling procedures are described in U.S. Pat. No. 4,302,438, which is hereby incorporated by reference. Iodinating, tritium labeling, and  $^{35}\text{S}$  labeling procedures especially adapted for murine monoclonal antibodies are described by Goding, J. W. (supra, pp 124-126) and the references cited therein, which are hereby incorporated by reference. Other procedures for iodinating biological agents, such as antibodies, binding portions thereof, probes, or ligands, are described by Hunter and Greenwood, *Nature* 144:945 (1962), David et al., *Biochemistry* 13:1014-1021 (1974), and U.S. Pat. Nos. 3,867,517 and 4,376,110, which are hereby incorporated by reference. Radiolabeling elements which are useful in imaging include  $^{123}\text{I}$ ,  $^{131}\text{I}$ ,  $^{111}\text{In}$ , and  $^{99\text{m}}\text{Tc}$ , for example. Procedures for iodinating biological agents are described by Greenwood, F. et al., *Biochem. J.* 89:114-123 (1963); Marchalonis, J., *Biochem. J.*

113:299-305 (1969); and Morrison, M. et al., *Immunochemistry*, 289-297 (1971), which are hereby incorporated by reference. Procedures for  $^{99\text{m}}\text{Tc}$ -labeling are described by Rhodes, B. et al. in Burchiel, S. et al. (eds.), *Tumor Imaging: The Radioimmunochemical Detection of Cancer*, New York: Masson 111-123 (1982) and the references cited therein, which are hereby incorporated by reference. Procedures suitable for  $^{111}\text{In}$ -labeling biological agents are described by Hnatowich, D. J. et al., *J. Immunol. Methods*, 65:147-157 (1983), Hnatowich, D. et al., *J. Applied Radiation*, 35:554-557 (1984), and Buckley, R. G. et al., *F.E.B.S.* 166:202-204 (1984), which are hereby incorporated by reference.

In the case of a radiolabeled biological agent, the biological agent is administered to the patient, is localized to the tumor bearing the antigen with which the biological agent reacts, and is detected or "imaged" in vivo using known techniques such as radionuclear scanning using e.g., a gamma camera or emission tomography. See e.g., A. R. Bradwell et al., "Developments in Antibody Imaging", *Monoclonal Antibodies for Cancer Detection and Therapy*, R. W. Baldwin et al., (eds.), pp. 65-85 (Academic Press 1985), which is hereby incorporated by reference. Alternatively, a positron emission transaxial tomography scanner, such as designated Pet VI located at Brookhaven National Laboratory, can be used where the radiolabel emits positrons (e.g.,  $^{11}\text{C}$ ,  $^{18}\text{F}$ ,  $^{15}\text{O}$ , and  $^{13}\text{N}$ ).

Fluorophore and chromophore labeled biological agents can be prepared from standard moieties known in the art. Since antibodies and other proteins absorb light having wavelengths up to about 310 nm, the fluorescent moieties should be selected to have substantial absorption at wavelengths above 310 nm and preferably above 400 nm. A variety of suitable fluorescers and chromophores are described by Stryer, *Science*, 162:526 (1968) and Brand, L. et al., *Annual Review of Biochemistry*, 41:843-868 (1972), which are hereby incorporated by reference. The biological agents can be labeled with fluorescent chromophore groups by conventional procedures such as those disclosed in U.S. Pat. Nos. 3,940,475, 4,289,747, and 4,376,110, which are hereby incorporated by reference.

One group of fluorescers having a number of the desirable properties described above are the xanthene dyes, which include the fluoresceins derived from 3,6-dihydroxy-9-henylxanthhydryl and resamines and rhodamines derived from 3,6-diamino-9-phenylxanthhydryl and lissanine rhodamine B. The rhodamine and fluorescein derivatives of 9-o-carboxyphenylxanthhydryl have a 9-o-carboxyphenyl group. Fluorescein compounds having reactive coupling groups such as amino and isothiocyanate groups such as fluorescein isothiocyanate and fluorescamine are readily available. Another group of fluorescent compounds are the naphthylamines, having an amino group in the  $\alpha$  or  $\beta$  position.

Biological agents can be labeled with fluorochromes or chromophores by the procedures described by Goding, J. (supra, pp 208-249). The biological agents can be labeled with an indicating group containing the NMR-active  $^{19}\text{F}$  atom, or a plurality of such atoms inasmuch as (i) substantially all of naturally abundant fluorine atoms are the  $^{19}\text{F}$  isotope and, thus, substantially all fluorine-containing compounds are NMR-active; (ii) many chemically active poly-fluorinated compounds such as trifluoroacetic anhydride are commercially available at relatively low cost, and (iii) many fluorinated compounds have been found medically acceptable for use in humans such as the perfluorinated polyethers utilized to carry oxygen as hemoglobin replacements. After

permitting such time for incubation, a whole body NMR determination is carried out using an apparatus such as one of those described by Pykett, *Scientific American*, 246:78-88 (1982), which is hereby incorporated by reference, to locate and image prostate epithelial cells.

In cases where it is important to distinguish between regions containing live and dead prostate epithelial cells or to distinguish between live and dead prostate epithelial cells, the antibodies of the present invention (or other biological agents of the present invention), labeled as described above, can be coadministered along with an antibody or other biological agent which recognizes only living or only dead prostate epithelial cells labeled with a label which can be distinguished from the label used to label the subject antibody. By monitoring the concentration of the two labels at various locations or times, spatial and temporal concentration variations of living and dead normal, benign hyperplastic, and cancerous prostate epithelial cells can be ascertained. In particular, this method can be carried out using the labeled antibodies of the present invention, which recognize both living and dead epithelial prostate cells, and labeled 7E11 antibodies, which recognize only dead epithelial prostate cells.

The biological agents can also be utilized to kill or ablate normal, benign hyperplastic, and cancerous prostate epithelial cells in vivo. This involves using the biological agents by themselves or with a cytotoxic drug to which the biological agents recognizing normal, benign hyperplastic, and cancerous prostate epithelial cells are bound. This involves administering the biological agents bonded to a cytotoxic drug to a mammal requiring such treatment. Since the biological agents recognize prostate epithelial cells, any such cells to which the biological agents bind are destroyed. Although such administration may destroy normal prostate epithelial cells, this is not problematic, because the prostate is not required for life or survival. Although the prostate may indirectly contribute to fertility, this is not likely to be a practical consideration in patients receiving the treatment of the present invention.

The biological agents of the present invention may be used to deliver a variety of cytotoxic drugs including therapeutic drugs, a compound emitting radiation, molecules of plants, fungal, or bacterial origin, biological proteins, and mixtures thereof. The cytotoxic drugs can be intracellularly acting cytotoxic drugs, such as short-range radiation emitters, including, for example, short-range, high-energy  $\alpha$ -emitters.

Enzymatically active toxins and fragments thereof are exemplified by diphtheria toxin A fragment, nonbinding active fragments of diphtheria toxin, exotoxin A (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain,  $\alpha$ -sacrin, certain *Aleurites fordii* proteins, certain Dianthin proteins, *Phytolacca americana* proteins (PAP, PAPII and PAP-S), *Morodica charantia* inhibitor, curcin, crotin, *Saponaria officinalis* inhibitor, gelonin, mitogillin, restrictocin, phenomycin, and enomycin, for example. Procedures for preparing enzymatically active polypeptides of the immunotoxins are described in W084/03508 and W085/03508, which are hereby incorporated by reference. Certain cytotoxic moieties are derived from adriamycin, chlorambucil, daunomycin, methotrexate, neocarzinostatin, and platinum, for example.

Procedures for conjugating the biological agents with the cytotoxic agents have been previously described. Procedures for conjugating chlorambucil with antibodies are described by Flechner, I., *European Journal of Cancer*, 9:741-745

(1973); Ghose, T. et al., *British Medical Journal*, 3:495-499 (1972); and Szekerke, M., et al., *Neoplasma*, 19:211-215 (1972), which are hereby incorporated by reference. Procedures for conjugating daunomycin and adriamycin to antibodies are described by Hurwitz, E. et al., *Cancer Research*, 35:1175-1181 (1975) and Arnon, R. et al. *Cancer Surveys*, 1:429-449 (1982), which are hereby incorporated by reference. Procedures for preparing antibody-ricin conjugates are described in U.S. Pat. No. 4,414,148 and by Osawa, T., et al. *Cancer Surveys*, 1:373-388 (1982) and the references cited therein, which are hereby incorporated by reference. Coupling procedures as also described in EP 86309516.2, which is hereby incorporated by reference.

In a particularly preferred embodiment of the present invention, a first biological agent is conjugated with a prodrug which is activated only when in close proximity with a prodrug activator. The prodrug activator is conjugated with a second biological agent according to the present invention, preferably one which binds to a non-competing site on the prostate specific membrane antigen molecule. Whether two biological agents bind to competing or non-competing binding sites can be determined by conventional competitive binding assays. For example, monoclonal antibodies J591, J533, and E99 bind to competing binding sites on the prostate specific membrane antigen molecule. Monoclonal antibody J415, on the other hand, binds to a binding site which is non-competing with the site to which J591, J533, and E99 bind. Thus, for example, the first biological agent can be one of J591, J533, and E99, and the second biological agent can be J415. Alternatively, the first biological agent can be J415, and the second biological agent can be one of J591, J533, and E99. Drug-prodrug pairs suitable for use in the practice of the present invention are described in Blakely et al., "ZD2767, an Improved System for Antibody-directed Enzyme Prodrug Therapy That Results in Tumor Regressions in Colorectal Tumor Xenografts," *Cancer Research*, 56:3287-3292 (1996), which is hereby incorporated by reference.

Alternatively, the biological agent can be coupled to high energy radiation emitters, for example, a radioisotope, such as  $^{131}\text{I}$ , a  $\gamma$ -emitter, which, when localized at the tumor site, results in a killing of several cell diameters. See, e.g., S. E. Order, "Analysis, Results, and Future Prospective of the Therapeutic Use of Radiolabeled Antibody in Cancer Therapy", *Monoclonal Antibodies for Cancer Detection and Therapy*, R. W. Baldwin et al. (eds.), pp 303-316 (Academic Press 1985), which is hereby incorporated by reference. Other suitable radioisotopes include  $\alpha$ -emitters, such as  $^{212}\text{Bi}$ ,  $^{213}\text{Bi}$ , and  $^{211}\text{At}$ , and  $\beta$ -emitters, such as  $^{186}\text{Re}$  and  $^{90}\text{Y}$ . Radiotherapy is expected to be particularly effective, because prostate cancer is a relatively radiosensitive tumor.

Where the biological agents are used alone to kill or ablate prostate epithelial cells, such killing or ablation can be effected by initiating endogenous host immune functions, such as complement-mediated or antibody-dependent cellular cytotoxicity.

The biological agent of the present invention can be used and sold together with equipment, as a kit, to detect the particular label.

Biological agents of the present invention can be used in conjunction with other therapeutic treatment modalities. Such other treatments include surgery, radiation, cryosurgery, thermotherapy, hormone treatment, chemotherapy, vaccines, and other immunotherapies.

Also encompassed by the present invention is a method of killing or ablating which involves using the biological

agents for prophylaxis. For example, these materials can be used to prevent or delay development or progression of prostate cancer.

Use of the prostate cancer therapy of the present invention has a number of benefits. Since the biological agents according to the present invention only target prostate epithelial cells, other tissue is spared. As a result, treatment with such biological agents is safer, particularly for elderly patients. Treatment according to the present invention is expected to be particularly effective, because it directs high levels of biological agents, such as antibodies or binding portions thereof, probes, or ligands, to the bone marrow and lymph nodes where prostate cancer metastases predominate. Moreover, tumor sites for prostate cancer tend to be small in size and, therefore, easily destroyed by cytotoxic agents. Treatment in accordance with the present invention can be effectively monitored with clinical parameters such as serum prostate specific antigen and/or pathological features of a patient's cancer, including stage, Gleason score, extracapsular, seminal, vesicle or perineural invasion, positive margins, involved lymph nodes, etc. Alternatively, these parameters can be used to indicate when such treatment should be employed.

Because the biological agents of the present invention bind to living prostate cells, therapeutic methods using these biological agents are much more effective than those which target lysed prostate cells. For the same reasons, diagnostic and imaging methods which determine the location of living normal, benign hyperplastic, or cancerous prostate epithelial cells are much improved by employing the biological agents of the present invention. In addition, the ability to differentiate between living and dead prostate cells can be advantageous, especially to monitor the effectiveness of a particular treatment regimen.

Hybridomas E99, J415, J533, and J591 have been deposited pursuant to, and in satisfaction of, the requirements of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure with the American Type Culture Collection ("A.T.C.C.") at 10801 University Boulevard, Manassas, Va. 20110. Hybridoma E99 was deposited on May 2, 1996, and received A.T.C.C. Designation Number HB-12101. Hybridoma J415 was deposited on May 30, 1996, and received A.T.C.C. Designation Number HB-12109. Hybridomas J533 and J591 were deposited on June 6, 1996, and received A.T.C.C. Designation Numbers HB-12127 and HB-12126, respectively.

The present invention is further illustrated by the following examples.

## EXAMPLES

### Example 1

#### Human Tissues

Fresh specimens of benign and malignant tissues were obtained from the Department of Pathology of New York Hospital Cornell University Medical Center ("NYH-CUMC").

### Example 2

#### Tissue Culture

Cultured cell lines of human cancers were obtained from the Laboratory of Urological Oncology of NYH-CUMC. The prostate cancer cell lines PC-3 (Mickey, D. D., et al., "Characterization Of A Human Prostate Adenocarcinoma Cell Line (DU145) As A Monolayer Culture And As A Solid Tumor In Athymic Mice," *Prog. Clin. Biol. Res.*, 37:67-84

(1980), which is hereby incorporated by reference), DU-145 (Mickey, D. D., et al., "Characterization Of A Human Prostate Adenocarcinoma Cell Line (DU145) As A Monolayer Culture And As A Solid Tumor In Athymic Mice," *Prog. Clin. Biol. Res.*, 37:67-84 (1980), which is hereby incorporated by reference), and LNCaP (Horoszewicz, J. S., et al., "LNCaP Model Of Human Prostatic Carcinoma," *Cancer Res.*, 43:1809-1818 (1983), which is hereby incorporated by reference) were obtained from the American Type Culture Collection (Rockville, Md.). Hybridomas were initially cloned in RPMI-1640 medium supplemented with 10% FCS, 0.1 mM nonessential amino acids, 2 mM L-glutamine, 100 units/ml of penicillin, 100 ug/ml of streptomycin and HAT medium (GIBCO, Grand Island, N.Y.). Subclones were cultured in the same medium without aminopterin.

### Example 3

#### Preparation of Mouse Monoclonal Antibodies

Female BALB/c mice were immunized intraperitoneally with LNCaP ( $6 \times 10^6$  cells) three times at 2 week intervals. A final intraperitoneal booster immunization was administered with fresh prostate epithelial cells which had been grown in vitro. Three days later, spleen cells were fused with SP-2 mouse myeloma cells utilizing standard techniques (Ueda, R., et al., "Cell Surface Antigens Of Human Renal Cancer Defined By Mouse Monoclonal Antibodies: Identification Of Tissue-Specific Kidney Glycoproteins," *Proc. Natl. Acad. Sci. USA*, 78:5122-5126 (1981), which is hereby incorporated by reference). Supernatants of the resulting clones were screened by rosette and complement cytotoxicity assays against viable LNCaP. Clones which were positive by these assays were screened by immunochemistry vs normal kidney, colon, and prostate. Clones which were LNCaP<sup>+</sup>/NmlKid<sup>-</sup>/colon<sup>-</sup>/prostate<sup>+</sup> were selected and subcloned 3 times by limiting dilution. The immunoglobulin class of cultured supernatant from each clone was determined by immunodiffusion using specified rabbit antisera (Calbiochem, San Diego, Calif.). mAbs were purified using the MAPS-II kit (Bio-Rad, Richmond, Calif.).

### Example 4

#### Biotinylation of mAbs

Purified mAbs were dialyzed in 0.1 M NaHCO<sub>3</sub> for 2 hours. One ml of mAb at 1 mg/ml was mixed with 0.1 ml of biotinamidocaproate N-hydroxysuccinamide ester (Sigma) in dimethylsulfoxide (1 mg/ml) and stirred for 4 hours at room temperature. Unbound biotin was removed by dialysis against phosphate buffered saline ("PBS").

### Example 5

#### Immunohistochemical Staining

Cryostat sections of prostate tissues were placed inside rings of Falcon 3034 plate covers (Becton-Dickenson, Lincoln Park, N.J.) previously coated with 0.45% gelatin solution as described in Marusich, M. F., "A Rapid Method For Processing Very Large Numbers Of Tissue Sections For Immunohistochemical Hybridoma Screening," *J. Immunol. Methods*, 111:143-145 (1988), which is hereby incorporated by reference. Plates were stored at -80° C. Cryostat sections were fixed with 2% paraformaldehyde in PBS for 10 min at room temperature, and, after washing with PBS, endogenous peroxidase activity was blocked by treatment with 0.3% hydrogen peroxide in PBS for 10 min at room temperature. After sections were incubated with 2% BSA in PBS for 20 min, mAbs were added for 60 min at room temperature. Slides were extensively washed with PBS and incubated with peroxidase-conjugated rabbit anti-mouse Ig (DAKO

Corp., Santa Barbara, Calif.) diluted 1:100 in 10% normal human serum in PBS for 60 min at room temperature. After a diaminobenzidine reaction, sections were counterstained with hematoxylin.

#### Example 6

##### Serological Analysis

The anti-mouse immunoglobulin mixed hemadsorption assay was performed as described in Ueda, R., et al., "Cell Surface Antigens Of Human Renal Cancer Defined By Mouse Monoclonal Antibodies: Identification Of Tissue-Specific Kidney Glycoproteins," *Proc. Natl. Acad. Sci. USA*, 78:5122-5126 (1981), which is hereby incorporated by reference. To prepare the indicator cells, anti-mouse Ig (DAKO Corp.) was conjugated to type O human RBC using 0.01% chromium chloride. Serological assays were performed on cells previously plated in Terasaki plates (Nunc, Denmark). Antibodies were incubated with target cells at room temperature for 1 hour. Target cells were then washed, and indicator cells added for 1 hour.

#### Example 7

##### Immunoprecipitation

LNCaP cells ( $2 \times 10^7$ ) were biotinylated with biotin-NHSS (at final concentration of 5 mM) for 30 minutes on ice. After washing, the biotinylated cells were resuspended in 1 ml lysis buffer (20 mM Tris/HCl pH 8.0, 1 mM EDTA, 1 mM PMSF, 1% triton X-100) for 30 min on ice. The suspension was centrifuged at 1500 g  $\times$  100 min at 4° C., and the supernatant was centrifuged at 12,000 rpm  $\times$  15 min at 4° C. The resulting lysate was preabsorbed with rabbit or goat anti-mouse IgG-coated pansorbin for 1 hour at 4° C. The pre-absorbed lysate was incubated with the mAb overnight at 4° C. Rabbit or goat anti-mouse Ig-coated agarose beads were added for 2 hours at 4° C. and then washed. The beads were resuspended in Tris-base/NaCl, added to sample buffer with 2-mercaptoethanol, and boiled for 5 min. After centrifuging, the supernatant was run on an SDS-PAGE 12% gel. The gel was transferred to a nitrocellulose membrane which was blocked and stained with streptavidin-peroxidase. The membrane was developed with diaminobenzidine ("DAB").

Sequential immunoprecipitation was similar except that the lysate was initially pre-cleared with one mAb overnight at 4° C. A second mAb was then used to immunoprecipitate the pre-cleared lysate.

Approximately 2000 clones were screened, of which four clones were selected as described in Example 3, above. After subcloning, supernatants from the 4 hybridomas, E99, J415, J533, and J591, were assayed by immunofluorescence against viable (i.e. unfixed) LNCaP, immunoprecipitation, and sequential immunoprecipitation to confirm reactivity to PSMA.

The immunofluorescence study using the LNCaP target cell (described originally in Horoszewicz, which is hereby incorporated by reference, to make the 7E11 antibody and the prototype cell line for expression for PSMA) shows that E99 antibody binds to and renders viable LNCaP cells immunofluorescent. This is in contrast to the 7E11 antibody, which, as noted originally in Horoszewicz, which is hereby incorporated by reference, gives only poor or no binding to viable LNCaP cells but exhibits strong binding once the cells are fixed (killed).

The reactivities of the four mAbs with normal human tissues were examined immunohistochemically; these results are presented in Table 3.

TABLE 3

Reactivity of mAbs with human normal tissues by indirect immunoperoxidase staining				
Tissues	E99 ( $\gamma_3$ )	J415 ( $\gamma_1$ )	J533 ( $\gamma_1$ )	J591 ( $\gamma_1$ )
Prostate*	●	●	●	●
Kidney				
Glomerulus	○	○	○	○
Prox. Tubule	■	■	■	■
Ureter	○	○	○	○
Bladder	○	○	○	○
Testis	○	○	○	○
Uterus	○			
Esophagus	○	○	○	○
Small Intestine	○	○	○	○
Stomach	○	○	○	○
Colon	○	○	○	○
Spleen	○	○	○	○
Thyroid	○	○	○	○
Lung	○	○	○	○
Pancreas	○	○	○	○
Liver	○	○	○	○
* BPH	0-3'	0-3'	0-4'	0-4'
* Prostate Cancer	0-3'	0-3'	0-4'	0-4'
* LNCaP (scid)	3'	3'	4'	4'
* LuCaP (scid)	0-2'	0-2'	0-3'	0-3'

● - positive; ■ - weak, heterogeneous; ○ - negative

The above sequential immunoprecipitation study showed that 7E11, E99, J415, J533, and J591 bind to the same molecule, i.e. PSMA.

#### Example 8

##### Western Blot Analysis

To confirm that antibodies E99, J415, J533, and J591 precipitate an identical band to the 7E11 antibody (i.e., PSMA), Western Blot analyses were performed. Seminal plasma (400  $\mu$ g/lane) or LNCaP lysate were loaded into lanes of 12% SDS-PAGE gels. After electrophoresis, the gels are transferred to nitrocellulose membranes. The membranes were blocked with 5% dry milk/Tris-buffered saline-tween 20 ("TBST") for 60 min at room temperature. After washing, the membranes were incubated with primary mAb for 60 min at room temperature. After repeat washing, the membranes were incubated with sheep anti-mouse-Ig-peroxidase 1/5000 in 5% dry milk/TBST for 60 min at room temperature. After repeat washing, the membranes were developed using a chemiluminescent tag designated "ECL" (Amersham Life Sciences, International, Arlington Heights, Ill.) according to the manufacturer's directions. The results of the Western Blot experiment are presented in Table 4.

TABLE 4

Western blot data					
Sample	7E11	E99	J415	J533	J591
Prostatic (seminal) fluid	100 KD band	100 KD band	100 KD band	100 KD band	100 KD band
LNCaP cell lysate	100 KD & 200 KD bands	100 KD & 200 KD bands	100 KD & 200 KD bands	100 KD & 200 KD bands	100 KD & 200 KD bands

#### Example 9

##### mAb Reactivity to External Domain of PSMA

To confirm cell surface (external) expression of the detected PSMA, fresh, viable LNCaP cells were tested, without fixation, in vitro, by immunofluorescence. LNCaP

cells were washed and incubated with mAb for 1 hour at room temperature and then with a rabbit anti-mouse Ig-fluorescein (DAKO Corp., Santa Barbara, Calif.). Wells were read with a fluorescent microscope. Negative control consisted of an isotype-matched irrelevant mAb, while an anti-class I MHC mAb served as a positive control.

Immunofluorescence and rosette assay results are presented in Table 5.

TABLE 5

Comparison of 7E11 with new mAbs					
LNCaP viable cells	7E11	E99	J415	J533	J591
Immunofluorescence	neg	3+	3+	4+	4+
Rosette assay	neg	+	+	+	+
LNCaP-fixed	+++	++++	+++	++	+++

## Example 10

## Competition Studies

A competition study was carried out to determine whether J591, J533, E99, and J415 detected the same or different antigenic sites (epitopes) of the prostate specific membrane antigen molecule using the following procedure.

Plates were coated with LNCaP cell line lysate as a source of prostate specific membrane antigen and washed to remove unbound material. "Cold" (unlabeled) monoclonal antibody was incubated on the plate for 1 hour at room temperature to allow binding to its antigenic site. Subsequently, a second monoclonal antibody, labeled either with biotin or <sup>125</sup>I, was added for an additional hour. Plates were washed to remove unbound material. The amount of the second monoclonal antibody bound to the prostate specific membrane antigen-coated plate was determined either by avidin-alkaline phosphatase in an enzyme-linked immunoassay (in the case of biotin-labeled second monoclonal antibody) or by physically counting the well in a gamma counter (in the case of <sup>125</sup>I-labeled second monoclonal antibody). Controls consisted of using the same monoclonal antibody both cold and labeled to define "100% competition" or using monoclonal antibody to a totally different molecule (e.g., monoclonal antibody I-56, which detects inhibin, a prostate related protein different from prostate specific membrane antigen) to define "0% competition".

The results indicated that J591, J533, and E99 each interfere, compete, or block binding of one another but do not block binding of J415 and vice versa. 7E11/CYT356, known to bind PSMA at a different (intracellular) site, did not block any of J591, J533, E99, or J415.

Having pairs of monoclonal antibodies which bind to non-competing sites permits development of antibody sandwich assays for detecting soluble antigens, such as solubilized prostate specific membrane antigen or fragment thereof, in, for example, body fluids. For example, the antigen (e.g., prostate specific membrane antigen or a fragment thereof) could be "captured" from body fluid with J591 and, in another step, detected by labeled J415.

In another setting, e.g. treatment, one could increase antibody binding by using a combination of non-competing monoclonal antibodies. For example, assuming the non-competing sites are each represented once on the prostate specific membrane antigen molecule, adding a combination of J591 plus J415 would bind twice as many monoclonal antibody molecules as either monoclonal antibody alone.

Binding two non-competing antigenic binding sites also can result in greater antigen cross-linking and, perhaps, increased internalization. Furthermore, since the two detected sites are physically located on the same prostate specific membrane antigen molecule, the binding of two monoclonal antibody molecules to that single prostate specific membrane antigen molecule puts the two monoclonal antibody molecules in close proximity to each other, a setting which provides optimal drug-prodrug interaction. For example, monoclonal antibody J591 can be conjugated with an inactive pro-drug and J415 can be conjugated with a pro-drug activator. Since prodrug and activator would be bound in close proximity only at the site of prostate specific membrane antigen-expressing cells (e.g., prostate cancer cells), prodrug activation to the active form would occur only at those sites.

## Example 11

## Microscopy

Confocal microscopy and immuno-electron microscopy demonstrated that E99, J591, J533, and J415 are bound to the cell membrane at clathrin-coated pits and then rapidly internalize into endosomes (cytoplasmic vesicles). FIGS. 1-4 are immuno-electron micrographs which follow the interaction of gold-labeled monoclonal antibody J591 with the cell surface as a function of time. In these figures, the location of the monoclonal antibody is indicated by the black dots.

Viable LNCaP cells were incubated with J591 for one hour at 4° C. The cells were washed and then held at 37° C. for 0, 5, 10, or 15 minutes, after which time they were fixed and processed for immuno-electron microscopy. FIG. 1 shows the cell prior to 37° C. incubation. J591 can be seen bound to the cell along the external aspect of the cell membrane. In this Figure, "M" denotes the cell's mitochondria, and "N" denotes its nucleus. FIG. 2 shows the cell after incubation at 37° C. for 5 minutes. The arrow indicates formation of a clathrin-coated pit. In FIG. 3, which shows the cell after a 10 minute 37° C. incubation, pinching off or endocytosis of the clathrin-coated pit can be seen, as indicated by the arrow. FIG. 4 shows that, after incubation at 37° C. for 15 minutes, monoclonal antibody J591 is contained in endocytic vesicles within the cell, as indicated by the arrows. As can be seen in FIG. 5, after incubation at 37° C. for 15 minutes, monoclonal antibody J591 is also contained within endosomes, as indicated by the arrows.

## Example 12

## Sequencing of the Variable Region of Monoclonal Antibody J591

Total RNA was prepared from 10<sup>7</sup> murine hybridoma J591 cells. A sample of the conditioned medium from these cells was tested for binding to the specific antigen for J591 on prostate cells. The conditioned medium was positive by both ELISA and Western Blot for binding to the antigen.

VH and VK cDNA were prepared using reverse transcriptase and mouse  $\kappa$  constant region and mouse IgG constant region primers. The first strand cDNAs were amplified by PCR using a variety of mouse signal sequence primers (6 for VH and 7 for VK). The amplified DNAs were gel-purified and cloned into the vector pT7Blue.

The VH and VK clones obtained were screened for correct inserts by PCR, and the DNA sequence of selected clones was determined by the dideoxy chain termination method.

Excluding the primer region (as the sequence of this depended on the sequence of the primer that was used), all the VH clones obtained gave identical sequence. This

sequence was obtained from clones produced with three different 5' primers. One clone had one base pair change within the signal sequence, and one clone contained an aberrant PCR product. Using the sequencing strategy shown in FIG. 6, the nucleotide sequence for the heavy chain was obtained. It is designated SEQ. ID. No. 1 and is presented in FIG. 7, along with the nucleotide sequence of the corresponding reverse, non-coding strand (designated SEQ. ID. No. 2). These sequences include part of the signal sequence and part of the constant region of the antibody. The corresponding deduced amino acid sequences of J591 VH, designated SEQ. ID. No. 3, SEQ. ID. No. 4, and SEQ. ID. No. 5, are also shown in FIG. 7. The coding strand of the J591 heavy chain's variable region (exclusive of signal sequence and constant region components) has the following nucleotide sequence (designated SEQ. ID. No. 6):

GAGGTCCAGCTGCAACAGTCTGGACCT-  
GAACTGGTGAAGCCTGGGACTTCAGTGAGG  
ATATCCTGCAAGACTTCTGGATACACAT-  
TCACTGAATATACCATACACTGGGTGAAG  
CAGAGCCATGGAAAGAGCCTTGAGTG-  
GATTGGAAACATCAATCCTAACAATGGTGGT  
ACCACCTACAATCAGAAGTTCGAGGAG-  
CAAGGCCACATTGACTGTAGACAAGTCTCC  
AGTACAGCCTACATGGAGCTCCGCAGC-  
CTAACATCTGAGGATTCTGCAGTCTATTAT TGTG-  
CAGCTGGTTGGAACCTTTGAC-  
TACTGGGCCAAGGCACCACTCTCACAGTCTCC  
TCA

The reverse, non-coding strand of the J591 heavy chain's variable region (exclusive of signal sequence and constant region components) has the following nucleotide sequence (designated SEQ. ID. No. 7):

TGAGGAGACTGTGAGAGTGGTGCCTTG-  
GCCCCAGTAGTCAAAGTTCCAACCAAGCTGC  
ACAATAATAGACTGCAGAATCCTCAGAT-  
GTTAGGCTGCGGAGCTCCATGTAGGCTGT ACTG-  
GAGGACTTGTCTACAGTCAATGTGGCCT-  
TGTCTCGAACTTCTGATTGTAGGT  
GGTACCACCATTTGTTAGGATTGAT-  
GTTTCCAATCCAATCAAGGCTCTTCCATGGCT  
CTGCTTCAACCCAGTGTATGGTATAT-  
TCAGTGAATGTGTATCCAGAAGTCTTGCAGGA  
TATCCTCACTGAAGTCCCAGGCTTTCAC-  
CAGTTCAGGTCCAGACTGTTGCAGCTGGAC CTC

The protein sequence corresponding to the J591 heavy chain's variable region (exclusive of signal sequence and constant region components) has the following nucleotide sequence (designated SEQ. ID. No. 8):

EVQLQSSGPPELVKPGTSLRISCKTSQY-  
FTFYTHWVKQSHGKSLWIGNINPNNGG TTYN-  
QKFEDKATLTVDKSSSTAYMELRSLT-  
SEDSAVVYCAAGWNFDYWGQGTILTVS S

The J591 VH is in Mouse Heavy Chains Subgroup IIA (designated SEQ. ID. No. 10) (Kabat et al., *Sequences of Proteins of Immunological Interest*, U.S. Department of Health and Human Services (1991) ("Kabat"), which is hereby incorporated by reference). The sequence of J591 VH is compared to the consensus sequence for this subgroup in FIG. 8.

In contrast to the VH, more than one VK sequence was obtained. Out of the 15 VK clones examined, four gave the sequence of an aberrant mouse Igκ from the fusion partner (Carol et al., *Molecular Immunology*, 25:991-995 (1988), which is hereby incorporated by reference). These clones originated from two specific 5' primers. No further work was done with these clones. Of the remaining clones, ten gave

identical nucleotide sequences, and one clone, VK17, gave an alternative VK sequence. The ten identical clones originated from three 5' primers (different from the two that gave the aberrant sequence), one of which also produced VK17. The sequencing strategy that was employed is shown in FIG. 9.

The nucleic acid sequence of J591 VK corresponding to the ten identical clones (designated SEQ. ID. No. 9) is presented in FIG. 10, along with the nucleic acid sequence of the corresponding reverse, non-coding strand (designated SEQ. ID. No. 10) and the deduced amino acid sequences, which are designated SEQ. ID. No. 11, SEQ. ID. No. 12, and SEQ. ID. No. 13. These sequences include part of the signal sequence and part of the constant region of the antibody. The coding strand of the J591 light (kappa) chain's variable region (exclusive of signal sequence and constant region components) corresponding to the ten identical clones has the following nucleotide sequence (designated SEQ. ID. No. 14):

AACATTGTAATGACCCAATCTC-  
CCAAATCCATGTCCATGTCAAGTAG-  
GAGAGAGGGTC ACCTTGACCTGCAAGGCCAGT-  
GAGAATGTGGTTACTTATGTTTCTGGTATCAACAG  
AAACCAGAGCAGTCTCTCTAAACTGCT-  
GATATACGGGGCATCCAACCGGTACACTGGG  
GTCCCCGATCGCTTCACAGGCAGTG-  
GATCTGCAACAGATTTCACTCTGACCATCAGC  
AGTGTGCAGGCTGAAGACCTTGCAGAT-  
TATCACTGTGGACAGGGTTACAGCTATCCG  
TACACGTTCCGGAGGGGGGACCAAGCTG-  
GAAATAAAA

The reverse, non-coding strand of the J591 light (kappa) chain's variable region (exclusive of signal sequence and constant region components) corresponding to the ten identical clones has the following nucleotide sequence (designated SEQ. ID. No. 15):

TTTTATTTCCAGCTTGGTCCCCCCTC-  
CGAACGTGTACGGATAGCTGTAACCCCTGTCC  
ACAGTGATAATCTGCAAGGCTCTTCAGC-  
CTGCACACTGCTGATGGTCAGAGTGAATC TGT-  
TGCAGATCCACTGCCTGTGAAGC-  
GATCGGGGACCCAGTGTACCGGTTGGATGC  
CCCGTATATCAGCAGTTTGGAGACT-  
GCTCTGGTTTCTGTTGATACCAGGAAACATA  
AGTAACCACATTCTCACTGGCCTTGCAG-  
GTCAAGGTGACCCCTCTCTCTACTGACAT GGA-  
CATGGATTGGGAGATTGGGTCAATACAATGTT

The protein sequence corresponding to the J591 light (kappa) chain's variable region (exclusive of signal sequence and constant region components) corresponding to the ten identical clones has the following nucleotide sequence (designated SEQ. ID. No. 16):

NIVMTQSPKMSMSVGERVTLTCKASEN-  
VVTYVSWYQQKPEQSPKLLIYGASNRYTG  
VPDRFTGSGSATDFTLTISSVQAEDLA-  
DYHCGQGYSPYTFGGGKLEIK

The coding strand of the J591 light (kappa) chain's variable region (exclusive of signal sequence and constant region components) corresponding to clone VK17 has the following nucleotide sequence (designated SEQ. ID. No. 17):

GACATTGTGATGACCCAGTCTCACAAAT-  
TCATGTCCACATCAGTAGGAGACAGGGTC  
AGCATCATCTGTAAGGCCAGTCAAGAT-  
GTGGGTACTGCTGTAGACTGGTATCAACAG  
AAACCAGGACAATCTCTAAACTACT-  
GATTTATTGGGCATCCACTCGGCACACTGGA



23

GTCCTGATCGCTTCACAGGCAGTG-  
GATCTGGGACAGACTTCACTCTCACCATTACT  
AATGTTTCAGTCTGAAGACTTGGCAGAT-  
TATTTCTGTCAGCAATATAACAGCTATCCT  
CTCACGTTTCGGTGCTGGGACCATGCTG- 5  
GACCTGAAA

The reverse, non-coding strand of the J591 light (kappa) chain's variable region (exclusive of signal sequence and constant region components) corresponding to clone VK17 has the following nucleotide sequence (designated SEQ. ID. No. 18):

TTTCAGGTCCAGCATGGTCCAGCAC-  
CGAACGTGAGAGGATAGCTGTTATATTGCTG  
ACAGAAATAATCTGCCAAGTCTTCA-  
GACTGAACATTAGTAATGGTGAGAGTGAAGTC 15  
TGTCCCAGATCCCACTGCCTGTGAAGC-  
GATCAGGGACTCCAGTGTGCCGAGTGGATGC  
CCAATAAATCAGTAGTTTAGGAGAT-  
TGTCCTGGTTTCTGTTGATACCACTCTACAGC  
AGTACCCACATCTTGACTGGCCTTACA- 20  
GATGATGCTGACCCTGTCTCTACTGATGT GGA-  
CATGAATTTGTGAGACTGGGTCATCACAATGTC

24

The protein sequence corresponding to the J591 light (kappa) chain's variable region (exclusive of signal sequence and constant region components) corresponding to clone VK17 has the following nucleotide sequence (designated SEQ. ID. No. 19):

DIVMTQSHKFMSTSVGDRVSIICKASQD-  
VGTAVDWYQQKPGQSPKLLIWASTRHTG  
VPDRFTGSGSGTDFTLTITNVQSEDLA-  
DYFCQQYNSYPLTFGAGTMLDLK

J591 VK is in the Mouse Kappa Chains Subgroup V (designated SEQ. ID. No. 21) (Kabat, which is hereby incorporated by reference). The sequence of J591 VK corresponding to the ten identical clones is compared to the consensus sequence for the subgroup in FIG. 11.

Although the invention has been described in detail for the purpose of illustration, it is understood that such detail is solely for that purpose and variations can be made by those skilled in the art without departing from the spirit and scope of the invention which is defined by the following claims.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 21

<210> SEQ ID NO 1

<211> LENGTH: 391

<212> TYPE: DNA

<213> ORGANISM: Mus sp.

<400> SEQUENCE: 1

tctcctgtca ggaactgcag gtgtcctctc tgagggtccag ctgcaacagt ctggacctga	60
actggtgaag cctgggactt cagtggagat atcctgcaag acttctggat acacattcac	120
tgaatatacc atacactggg tgaagcagag ccatggaaag agccttgagt ggattggaaa	180
catcaatcct aacaatgggt gtaccaccta caatcagaag ttcgaggaca aggccacatt	240
gactgtagac aagtctcca gtacagccta catggagctc cgcagcctaa catctgagga	300
ttctgcagtc tattattgtg cagctggttg gaactttgac tactggggcc aaggcaccac	360
tctcacagtc tctcagcca aaacgacacc c	391

<210> SEQ ID NO 2

<211> LENGTH: 391

<212> TYPE: DNA

<213> ORGANISM: Mus sp.

<400> SEQUENCE: 2

gggtgtcgtt ttggctgagg agactgtgag agtgggtcct tggccccagt agtcaaagtt	60
ccaaccagct gcacaataat agactgcaga atcctcagat gttaggctgc ggagctccat	120
gtaggctgta ctggaggact tgtctacagt caatgtggcc ttgtcctcga acttctgatt	180
gtagggtgta ccaccattgt taggattgat gtttccaatc cactcaaggc tctttccatg	240
gctctgcttc acccagtgta tggatatatt agtgaatgtg tatccagaag tcttgcagga	300
tatcctcact gaagtcacag gcttcaccag ttcagggtcca gactgttgca gctggacctc	360
agagaggaca cctgcagttc ctgacaggag a	391

<210> SEQ ID NO 3

-continued

<211> LENGTH: 123  
 <212> TYPE: PRT  
 <213> ORGANISM: Mus sp.

<400> SEQUENCE: 3

```

Ser Pro Val Arg Asn Cys Arg Cys Pro Leu Gly Pro Ala Ala Thr Val
1           5           10           15
Trp Thr Thr Gly Glu Ala Trp Asp Phe Ser Glu Asp Ile Leu Gln Asp
20           25           30
Phe Trp Ile His Ile His Ile Tyr His Thr Leu Gly Glu Ala Glu Pro
35           40           45
Trp Lys Glu Pro Val Asp Trp Lys His Gln Ser Gln Trp Trp Tyr His
50           55           60
Leu Gln Ser Glu Val Arg Gly Gln Gly His Ile Asp Cys Arg Gln Val
65           70           75           80
Leu Gln Tyr Ser Leu His Gly Ala Pro Gln Pro Asn Ile Gly Phe Cys
85           90           95
Ser Leu Leu Leu Cys Ser Trp Leu Glu Leu Leu Gly Pro Arg His
100          105          110
His Ser His Ser Leu Leu Ser Gln Asn Asp Thr
115          120

```

<210> SEQ ID NO 4  
 <211> LENGTH: 130  
 <212> TYPE: PRT  
 <213> ORGANISM: Mus sp.

<400> SEQUENCE: 4

```

Leu Leu Ser Gly Thr Ala Gly Val Leu Ser Glu Val Gln Leu Gln Gln
1           5           10           15
Ser Gly Pro Glu Leu Val Lys Pro Gly Thr Ser Val Arg Ile Ser Cys
20           25           30
Lys Thr Ser Gly Tyr Thr Phe Thr Glu Tyr Thr Ile His Trp Val Lys
35           40           45
Gln Ser His Gly Lys Ser Leu Glu Trp Ile Gly Asn Ile Asn Pro Asn
50           55           60
Asn Gly Gly Thr Thr Tyr Asn Gln Lys Phe Glu Asp Lys Ala Thr Leu
65           70           75           80
Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr Met Glu Leu Arg Ser Leu
85           90           95
Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys Ala Ala Gly Trp Asn Phe
100          105          110
Asp Tyr Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser Ala Lys Thr
115          120          125
Thr Pro
130

```

<210> SEQ ID NO 5  
 <211> LENGTH: 125  
 <212> TYPE: PRT  
 <213> ORGANISM: Mus sp.

<400> SEQUENCE: 5

```

Leu Ser Cys Gln Glu Leu Gln Val Ser Ser Leu Arg Ser Ser Cys Asn
1           5           10           15
Ser Leu Asp Leu Asn Trp Ser Leu Gly Leu Gln Gly Tyr Pro Ala Arg
20           25           30

```

-continued

Leu Leu Asp Thr His Ser Leu Asn Ile Pro Tyr Thr Gly Ser Arg Ala  
 35 40 45  
 Met Glu Arg Ala Leu Ser Gly Leu Glu Thr Ser Ile Leu Thr Met Val  
 50 55 60  
 Val Pro Pro Thr Ile Arg Ser Ser Arg Thr Arg Pro His Leu Thr Ser  
 65 70 75 80  
 Pro Pro Val Gln Pro Thr Trp Ser Ser Ala Ala His Leu Arg Ile Leu  
 85 90 95  
 Gln Ser Ile Ile Val Gln Leu Val Gly Thr Leu Thr Thr Gly Ala Lys  
 100 105 110  
 Ala Pro Leu Ser Gln Pro Ser Gln Pro Lys Arg His Pro  
 115 120 125

<210> SEQ ID NO 6  
 <211> LENGTH: 345  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus sp.

<400> SEQUENCE: 6

gaggtccagc tgcaacagtc tggacctgaa ctggtgaagc ctgggacttc agtgaggata 60  
 tcctgcaaga cttctggata cacattcact gaatatacca tacactgggt gaagcagagc 120  
 catggaaaga gccttgatgt gattggaaac atcaatccta acaatgggtg taccacctac 180  
 aatcagaagt tcgaggacaa ggccacattg actgtagaca agtctccag tacagcctac 240  
 atggagctcc gcagcctaac atctgaggat tctgcagtct attattgtgc agctggttgg 300  
 aactttgact actggggcca aggcaccact ctccacagtct cctca 345

<210> SEQ ID NO 7  
 <211> LENGTH: 345  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus sp.

<400> SEQUENCE: 7

tgaggagact gtgagagtgg tgccttgcc ccagtagtca aagttccaac cagctgcaca 60  
 ataatagact gcagaatcct cagatgttag gctgcggagc tccatgtagg ctgtactgga 120  
 ggacttgtct acagtcaatg tggccttgct ctgaacttc tgattgtagg tggaccacc 180  
 attgttagga ttgatgttc caatccactc aaggtctctt ccatggctct gcttcaccca 240  
 gtgtatgta tattcagtga atgtgtatcc agaagtcttg caggatatcc tcaactgaagt 300  
 cccaggttc accagttcag gtccagactg ttgcagctgg acctc 345

<210> SEQ ID NO 8  
 <211> LENGTH: 115  
 <212> TYPE: PRT  
 <213> ORGANISM: Mus sp.

<400> SEQUENCE: 8

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Thr  
 1 5 10 15  
 Ser Val Arg Ile Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Glu Tyr  
 20 25 30  
 Thr Ile His Trp Val Lys Gln Ser His Gly Lys Ser Leu Glu Trp Ile  
 35 40 45  
 Gly Asn Ile Asn Pro Asn Asn Gly Gly Thr Thr Tyr Asn Gln Lys Phe  
 50 55 60  
 Glu Asp Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr

-continued

65	70	75	80	
Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys				
	85	90	95	
Ala Ala Gly Trp Asn Phe Asp Tyr Trp Gly Gln Gly Thr Thr Leu Thr				
	100	105	110	
Val Ser Ser				
	115			

<210> SEQ ID NO 9  
 <211> LENGTH: 363  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus sp.  
 <400> SEQUENCE: 9

ttatatggag ctgatgggaa cattgtaatg acccaatctc ccaaatccat gtccatgtca	60
gtaggagaga ggggtcacctt gacctgcaag gccagtgaga atgtggttac ttatgtttcc	120
tggtatcaac agaaaccaga gcagtctcct aaactgtga tatacggggc atccaaccgg	180
tacactgggg tccccgatcg cttcacaggc agtggatctg caacagattt cactctgacc	240
atcagcagtg tgcaggctga agaccttgca gattatcact gtggacaggg ttacagctat	300
ccgtacacgt tcggaggggg gaccaagctg gaaataaaac gggctgatgc tgcaccaact	360
gta	363

<210> SEQ ID NO 10  
 <211> LENGTH: 363  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus sp.  
 <400> SEQUENCE: 10

tacagttggt gcagcatcag cccgttttat ttccagcttg gtccccctc cgaacgtgta	60
cggtagatctg taacctgtc cacagtgaata atctgcaagg tcttcagcct gcacactgct	120
gatggtcaga gtgaaatctg ttgcagatcc actgcctgtg aagcgatcgg ggaccccgat	180
gtaccggttg gatgcccctg atatcagcag tttaggagac tgctctggtt tctgttgata	240
ccaggaaaca taagtaacca cattctcact ggccctgcag gtcaagtgta cccctctctc	300
tactgacatg gacatggatt tgggagattg ggtcattaca atgttcccat cagctccata	360
taa	363

<210> SEQ ID NO 11  
 <211> LENGTH: 121  
 <212> TYPE: PRT  
 <213> ORGANISM: Mus sp.  
 <400> SEQUENCE: 11

Leu Tyr Gly Ala Asp Gly Asn Ile Val Met Thr Gln Ser Pro Lys Ser	
1	15
Met Ser Met Ser Val Gly Glu Arg Val Thr Leu Thr Cys Lys Ala Ser	
20	30
Glu Asn Val Val Thr Tyr Val Ser Trp Tyr Gln Gln Lys Pro Glu Gln	
35	45
Ser Pro Lys Leu Leu Ile Tyr Gly Ala Ser Asn Arg Tyr Thr Gly Val	
50	60
Pro Asp Arg Phe Thr Gly Ser Gly Ser Ala Thr Asp Phe Thr Leu Thr	
65	80
Ile Ser Ser Val Gln Ala Glu Asp Leu Ala Asp Tyr His Cys Gly Gln	

-continued

				85						90									95
Gly	Tyr	Ser	Tyr	Pro	Tyr	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile				
				100				105						110					
Lys	Arg	Ala	Asp	Ala	Ala	Pro	Thr	Val											
		115						120											

<210> SEQ ID NO 12  
 <211> LENGTH: 114  
 <212> TYPE: PRT  
 <213> ORGANISM: Mus sp.

<400> SEQUENCE: 12

Tyr	Met	Glu	Leu	Met	Gly	Thr	Leu	Pro	Asn	Leu	Pro	Asn	Pro	Cys	Pro				
1				5					10					15					
Cys	Gln	Glu	Arg	Gly	Ser	Pro	Pro	Ala	Arg	Pro	Val	Arg	Met	Trp	Leu				
			20					25					30						
Leu	Met	Phe	Pro	Gly	Ile	Asn	Arg	Asn	Gln	Ser	Ser	Leu	Leu	Asn	Cys				
		35				40						45							
Tyr	Thr	Gly	His	Pro	Thr	Gly	Thr	Leu	Gly	Ser	Pro	Ile	Ala	Ser	Gln				
		50				55					60								
Ala	Val	Asp	Leu	Gln	Gln	Ile	Ser	Leu	Pro	Ser	Ala	Val	Cys	Arg	Leu				
65				70					75					80					
Lys	Thr	Leu	Gln	Ile	Ile	Thr	Val	Asp	Arg	Val	Thr	Ala	Ile	Arg	Thr				
			85					90						95					
Arg	Ser	Glu	Gly	Gly	Pro	Ser	Trp	Lys	Asn	Gly	Leu	Met	Leu	His	Gln				
			100					105					110						

Leu Tyr

<210> SEQ ID NO 13  
 <211> LENGTH: 116  
 <212> TYPE: PRT  
 <213> ORGANISM: Mus sp.

<400> SEQUENCE: 13

Ile	Ile	Trp	Ser	Trp	Glu	His	Cys	Asn	Asp	Pro	Ile	Ser	Gln	Ile	His				
1				5				10						15					
Val	His	Val	Ser	Arg	Arg	Glu	Gly	His	Leu	Asp	Leu	Gln	Gly	Gln	Glu				
			20					25					30						
Cys	Gly	Tyr	Leu	Cys	Phe	Leu	Val	Ser	Thr	Glu	Thr	Arg	Ala	Val	Ser				
			35				40					45							
Thr	Ala	Asp	Ile	Arg	Gly	Ile	Gln	Pro	Val	His	Trp	Gly	Pro	Arg	Ser				
		50				55					60								
Leu	His	Arg	Gln	Trp	Ile	Cys	Asn	Arg	Phe	His	Ser	Asp	His	Gln	Gln				
65				70				75						80					
Cys	Ala	Gly	Arg	Pro	Cys	Arg	Leu	Ser	Leu	Trp	Thr	Gly	Leu	Gln	Leu				
			85					90						95					
Ser	Val	His	Val	Arg	Arg	Gly	Asp	Gln	Ala	Gly	Asn	Lys	Thr	Gly	Cys				
			100				105						110						

Cys Thr Asn Cys  
115

<210> SEQ ID NO 14  
 <211> LENGTH: 321  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus sp.

<400> SEQUENCE: 14

-continued

---

```

aacattgttaa tgaccaaatc tcccaaatcc atgtccatgt cagtaggaga gagggtcacc 60
ttgacctgca aggccagtga gaatgtggtt acttatgttt cctggatatca acagaaacca 120
gagcagctctc cttaaactgct gatatacggg gcatccaacc ggtacactgg ggtccccgat 180
cgcttcacag gcagtggatc tgcaacagat ttcactctga ccatcagcag tgtgcaggct 240
gaagaccttg cagattatca ctgtggacag ggttacagct atccgtacac gttcggaggg 300
gggaccaagc tggaaataaa a 321

```

```

<210> SEQ ID NO 15
<211> LENGTH: 321
<212> TYPE: DNA
<213> ORGANISM: Mus sp.

```

```

<400> SEQUENCE: 15

```

```

ttttatttcc agcttgggcc cccctccgaa cgtgtacgga tagctgtaac cctgtccaca 60
gtgataatct gcaaggtctt cagcctgcac actgctgatg gtcagagtga aatctgttgc 120
agatccactg cctgtgaagc gatcggggac cccagtgtac cggttggatg ccccgatat 180
cagcagttta ggagactgct ctggtttctg ttgataccag gaaacataag taaccacatt 240
ctcactggcc ttgcaggta aggtgaccct ctctcctact gacatggaca tggatttggg 300
agattgggtc attacaatgt t 321

```

```

<210> SEQ ID NO 16
<211> LENGTH: 107
<212> TYPE: PRT
<213> ORGANISM: Mus sp.

```

```

<400> SEQUENCE: 16

```

```

Asn Ile Val Met Thr Gln Ser Pro Lys Ser Met Ser Met Ser Val Gly
1           5           10          15
Glu Arg Val Thr Leu Thr Cys Lys Ala Ser Glu Asn Val Val Thr Tyr
20          25          30
Val Ser Trp Tyr Gln Gln Lys Pro Glu Gln Ser Pro Lys Leu Leu Ile
35          40          45
Tyr Gly Ala Ser Asn Arg Tyr Thr Gly Val Pro Asp Arg Phe Thr Gly
50          55          60
Ser Gly Ser Ala Thr Asp Phe Thr Leu Thr Ile Ser Ser Val Gln Ala
65          70          75          80
Glu Asp Leu Ala Asp Tyr His Cys Gly Gln Gly Tyr Ser Tyr Pro Tyr
85          90          95
Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100         105

```

```

<210> SEQ ID NO 17
<211> LENGTH: 321
<212> TYPE: DNA
<213> ORGANISM: Mus sp.

```

```

<400> SEQUENCE: 17

```

```

gacattgtga tgaccacgct tcacaaatcc atgtccacat cagtaggaga cagggtcagc 60
atcatctgta aggccagtca agatgtgggt actgctgtag actggtatca acagaaacca 120
ggacaatctc cttaaactact gatttattgg gcatccactc ggcacactgg agtccctgat 180
cgcttcacag gcagtggatc tgggacagac ttcactctca ccattactaa tgttcagtct 240
gaagacttgg cagattattt ctgtcagcaa tataacagct atcctctcac gttcgggtgt 300

```

-continued

gggacccatgc tggacctgaa a

321

<210> SEQ ID NO 18  
 <211> LENGTH: 321  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus sp.

&lt;400&gt; SEQUENCE: 18

tttcagggtcc agcatgggtcc cagcaccgaa cgtgagagga tagctgttat attgctgaca	60
gaaataatct gccaaagtctt cagactgaac attagtaatg gtgagagtga agtctgtccc	120
agatccactg cctgtgaagc gatcaggac tccagtgtgc cgagtggatg cccaataaat	180
cagtagttta ggagattgtc ctggtttctg ttgataccag tctacagcag taccacatc	240
ttgactggcc ttacagatga tgctgacct gtctcctact gatgtggaca tgaatttg	300
agactgggtc atcacaatgt c	321

<210> SEQ ID NO 19  
 <211> LENGTH: 107  
 <212> TYPE: PRT  
 <213> ORGANISM: Mus sp.

&lt;400&gt; SEQUENCE: 19

Asp Ile Val Met Thr Gln Ser His Lys Phe Met Ser Thr Ser Val Gly	
1 5 10 15	
Asp Arg Val Ser Ile Ile Cys Lys Ala Ser Gln Asp Val Gly Thr Ala	
20 25 30	
Val Asp Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile	
35 40 45	
Tyr Trp Ala Ser Thr Arg His Thr Gly Val Pro Asp Arg Phe Thr Gly	
50 55 60	
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Thr Asn Val Gln Ser	
65 70 75 80	
Glu Asp Leu Ala Asp Tyr Phe Cys Gln Gln Tyr Asn Ser Tyr Pro Leu	
85 90 95	
Thr Phe Gly Ala Gly Thr Met Leu Asp Leu Lys	
100 105	

<210> SEQ ID NO 20  
 <211> LENGTH: 125  
 <212> TYPE: PRT  
 <213> ORGANISM: Mus sp.

&lt;400&gt; SEQUENCE: 20

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala	
1 5 10 15	
Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr	
20 25 30	
Tyr Met Asn Asn Trp Val Lys Gln Ser Pro Gly Lys Ser Leu Glu Trp	
35 40 45	
Ile Gly Asp Ile Asn Pro Gly Asn Gly Gly Thr Ser Tyr Asn Gln Lys	
50 55 60	
Phe Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala	
65 70 75 80	
Tyr Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr	
85 90 95	
Cys Ala Arg Gly Tyr Tyr Ser Ser Ser Tyr Met Ala Tyr Tyr Ala Phe	
100 105 110	

-continued

Asp Tyr Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
115 120 125

<210> SEQ ID NO 21  
<211> LENGTH: 109  
<212> TYPE: PRT  
<213> ORGANISM: Mus sp.

<400> SEQUENCE: 21

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Leu Gly  
1 5 10 15  
Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Asp Asp Ile Ser Asn  
20 25 30  
Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Gly Ser Pro Lys Leu Leu  
35 40 45  
Ile Tyr Tyr Ala Ser Arg Leu His Ser Gly Val Pro Ser Arg Phe Ser  
50 55 60  
Gly Ser Gly Ser Gly Thr Asp Tyr Ser Leu Thr Ile Ser Asn Leu Glu  
65 70 75 80  
Gln Glu Asp Ile Ala Thr Tyr Phe Cys Gln Gln Gly Asn Thr Leu Pro  
85 90 95  
Pro Arg Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys  
100 105

What is claimed:

1. A method of ablating or killing normal, benign hyperplastic prostate epithelial cells or cancerous prostate epithelial cells in a patient comprising administering:

(a) an isolated antibody or antigen binding portion thereof which binds to the extracellular domain of prostate specific membrane antigen (PSMA) which binding occurs on living cells;

contacting said cells with the isolated antibody or antigen binding portion thereof under conditions effective to permit binding of the isolated antibody or antigen binding portion thereof to the extracellular domain of the PSMA wherein immune effector cells and/or complement components are recruited to the antibody/cell complex, thereby ablating or killing said cells and administering;

(b) a second isolated antibody or antigen binding portion thereof which binds to the extracellular domain of PSMA; contacting said cells with the second isolated antibody or antigen binding portion thereof under conditions effective to permit binding of the second isolated antibody or antigen binding portion to the extra-

cellular domain of the PSMA wherein immune effector cells and/or complement components are recruited to the antibody/cell complex, thereby ablating or killing said cells.

2. A method according to claim 1, wherein the isolated antibody or antigen binding portion thereof and the second isolated antibody or antigen binding portion thereof bind to non-competing binding sites on the extracellular domain of the PSMA.

3. A method according to claim 1, wherein the isolated antibody or antigen binding portion thereof is a monoclonal antibody or antigen binding portion thereof wherein the monoclonal antibody is produced by a hybridoma having an ATCC accession Number of HB-12126 and the second isolated antibody or antigen binding portion thereof is a monoclonal antibody or antigen binding portion thereof, wherein the second antibody is a monoclonal antibody produced by a hybridoma having an ATCC Accession Number selected from the group consisting of HB-12101, HB-12109, and HB-12127.

\* \* \* \* \*